

# Comparison of White Crappie Populations in Diked and Undiked Lake Erie Wetlands<sup>1</sup>

CAROL A. MARKHAM, WILLIAM E. LYNCH JR., DAVID L. JOHNSON, AND RAYMOND W. PETERING, School of Natural Resources, The Ohio State University, 2021 Coffey Road, Columbus, OH 43210

**ABSTRACT.** Most of Ohio's remaining Lake Erie wetlands are diked to enhance habitat diversity. There is concern that fish communities in these wetlands may be isolated from adjacent waters. However, little data are available with which to evaluate possible isolation. We conducted a study that examined spring length frequencies, age structure, and growth of white crappie (*Pomoxis annularis*) populations in 3 diked wetlands and 2 undiked, adjacent areas. If populations are not isolated then differences in population parameters between the two types of systems should not be evident. White crappies were collected in April-May 1987 using trap nets. Length frequencies of white crappies were not similar between diked wetlands and adjacent areas, and mean lengths were significantly less in the diked wetlands. Populations were not comprised of similar age classes in the 5 systems. White crappies in diked wetlands grew significantly slower than their conspecifics in the undiked areas. These data indicate that white crappies in diked wetlands are isolated from populations in undiked areas even though up to 75% of water in diked wetlands can be exchanged each year.

OHIO J. SCI. 97 (4): 72-77, 1997

## INTRODUCTION

Wetlands and their diverse macrophyte communities provide important habitats for a variety of fish species (Liston and Chubb 1985, Johnson 1989). Extensive agricultural, commercial, and industrial pressures have reduced these coastal wetlands along the western basin of Lake Erie from about 4,000 km<sup>2</sup> to about 100 km<sup>2</sup> (Herdendorf 1987). Most (84%) of the remaining 100 km<sup>2</sup> has been diked by both private and governmental entities so that water level manipulations can create enhanced wildlife habitats (Bookhout and others 1989). The few remaining undiked areas provide little in the way of quality habitat as the effects of carp, wind, water level fluctuations, and agriculture have combined to create very turbid systems containing little or no vegetation.

Regulatory agencies have become concerned about the impact of diking on fish communities. Of special concern is whether the dikes are restricting fish movement into and out of wetlands containing most of the remaining quality wetland habitat in western Lake Erie. Johnson and others (1997), in a study of a diked wetland and its adjacent undiked area prior to ours, concluded that the fish community in the diked wetland appeared to be isolated. Catch rates, sizes collected, and body condition of species common to both areas were significantly different between these areas. Subpopulations of a fish species can result when part of a population becomes spatially separated from the remainder of that population. If isolation is of sufficient duration, these subpopulations may begin to show differences in population parameters such as age structure, growth rates, and condition (Everhart and others 1975).

In this paper, we present length frequencies, age structures, and growth rate data of white crappie

(*Pomoxis annularis*) collected from 3 diked wetlands and 2 adjacent undiked areas of Lake Erie. Comparison of these parameters will allow us to evaluate if populations of white crappie in diked wetlands are likely isolated from adjacent undiked areas with respect to population form. In addition, these data will provide additional insight into the impacts of diking and management strategies on fish communities.

## MATERIAL AND METHODS

### Study Areas

White crappie were collected from 3 diked wetlands, two of which were located in western Sandusky Bay with the remaining diked wetland being adjacent to Lake Erie. Muddy Creek Bay and Crane Creek were the undiked areas sampled along Sandusky Bay and Lake Erie, respectively. Rusk #2 (hereafter Rusk) was an 18 ha privately-owned diked wetland on the north shore of Sandusky Bay and receives its water via a township canal connected to Sandusky Bay. Boardman's Unit (hereafter OSC) was a 40 ha diked wetland within the Ottawa Shooting Club on the south side of Muddy Creek Bay. Its water source was Muddy Creek Bay. The remaining diked wetland sampled was pool 2b (hereafter ONWR) on the Ottawa National Wildlife Refuge, a 38 ha wetland receiving its water from Crane Creek near its confluence with Lake Erie. All 3 wetlands contained diverse macrophyte communities. Muddy Creek Bay and the Crane Creek Embayment are large shallow, undiked areas (>100 ha) characterized by turbid water and little vegetation.

### Fish Collection

Juvenile and adult white crappie were collected from 10 April through 17 May 1987. Spring sampling was done to insure that yearly growth annuli had not yet been laid down. Five South Dakota style trap nets (1.5 m square, 30-m leads, 13 mm mesh) were fished about

<sup>1</sup>Manuscript received 21 February 1997 and in revised form 14 August 1997 (#97-04).

21 h per day for 2-4 d at each of the 5 areas. Trap net locations within each study area were randomly selected at the beginning of each sampling period. White crappies collected were measured to the nearest millimeter (TL, total length). For each area, a subsample of 5 white crappie per 10 mm length group (range 50-350+ mm TL) were frozen for later growth work.

### Length Frequencies

Length frequencies (25 mm groups) were constructed and Kolmogorov-Smirnov's (K-S) test (Hollander and Wolfe 1973) was used to determine if white crappie length distributions were different between diked and undiked areas within the Sandusky Bay or Lake Erie watersheds. A proportional stock density (PSD) was calculated for each wetland. PSD is a numerical descriptor of length frequency data and can provide insight into population structure and dynamics (Anderson and Neumann 1996). PSDs were calculated as follows:

$$\text{PSD} = (\# \text{ of fish } > 200 \text{ mm} / \# \text{ of fish } \geq 130 \text{ mm}) * 100$$

### Age and Growth

We chose to use otoliths for determining age and growth rather than scales because of greater accuracy associated with otoliths (Boxrucker 1986). The assumption that otolith radius is proportional to total fish length has been validated (Boxrucker 1986, Schramm and Doerzbacher 1982). Otoliths were removed and placed in glycerine for 1-2 weeks to clear and then examined using methods described by Schramm and Doerzbacher (1982). Each otolith was examined by 2 individuals and the age of the fish was accepted once both individuals agreed. If agreement could not be reached, the otolith was discarded from the data set. Length-at-age (growth) for each fish was calculated as follows:

$$Li = (ABi / OR) * TL$$

where  $Li$  = length at age  $i$ ,  $ABi$  = distance to the annual band associated with age  $i$ ,  $OR$  = otolith radius, and  $TL$  = total fish length. From these data, mean lengths-at-age were calculated for each age class at each of the 5 sample areas. Analysis of variance (ANOVA) was used to statistically compare these means between areas within a watershed. Age frequencies were constructed and Kolmogorov-Smirnov's (K-S) test was used to determine if white crappie age distributions were different between diked and undiked areas.

## RESULTS

A total of 1031 trap-net hours resulted in a catch of 239 white crappies, all areas combined. Catches per 24-h period in Muddy Creek Bay and the Crane Creek Embayment were 7.6 and 3.4 fish, respectively. Catches in OSC, ONWR, and Rusk wetlands were 3.8, 4.8, and 11.0 fish per 24 h, respectively.

### Length Frequencies

Length frequencies of white crappies varied considerably among the 5 areas sampled (Fig. 1). Within both the Sandusky Bay and Lake Erie watersheds, length

distributions were dissimilar between the diked wetlands and the adjacent undiked area (K-S test,  $P < 0.05$ ). PSDs for the 2 undiked areas (100 & 96) exceeded values from the diked areas in both watersheds. Catches in undiked areas were composed largely of adult white crappie ( $> 225$  mm) whereas catches in diked wetlands contained juveniles and adults with the exception of Rusk (Fig. 1).

### Age and Growth

Age distributions of white crappies varied considerably among the 5 areas (Fig. 2). Within both watersheds, age distributions were dissimilar between the diked wetlands and the adjacent undiked area (K-S test,  $P < 0.05$ ).

Average length-at-age (all year classes combined) was less in diked wetlands than in the adjacent undiked areas for both the Sandusky Bay and Lake Erie areas (Fig. 3). Comparisons of length-at-age for the age classes (2-4) common to the systems indicated that white crappie growth was significantly less in the diked wetlands as compared to undiked areas (Table 1, ANOVA,  $P < 0.05$ ). In undiked areas, white crappie grew quickly to 270 and 297 mm TL on average in Crane Creek and Muddy Creek Bay by age 4. Length-at-age varied considerably among the diked wetlands at all ages with Rusk on Sandusky Bay exhibiting very slow white crappie growth while OSC exhibited the fastest growth. Four year old fish averaged 187 mm TL in Rusk and 261 mm TL in OSC.

## DISCUSSION

Johnson and others (1997) provided the first evidence that fish communities in diked wetlands may be biologically isolated from communities in adjacent undiked areas. Their working premise was that considerable exchange of fish or similar conditions within diked and undiked areas would create similar population parameters for each species. However, they found that mean lengths of 5 fish species common to both a diked and undiked area were significantly smaller in the diked wetland. Body condition factors of white crappie and brown bullhead were also lower for populations in the diked wetland. These data suggest that fish communities in their diked wetland were probably isolated from adjacent Muddy Creek Bay. They also noted that ecological conditions between the two areas are different and can cause biological isolation even though fish exchange may occur. Johnson and others (1997) noted that growth information was needed and additional wetlands needed to be sampled before it can be conclusively decided that diked wetland fish communities are isolated.

Our study of white crappie provides additional evidence that fish communities in diked wetland may be biologically isolated. In both the Lake Erie and Sandusky Bay watersheds, we found that spring length frequencies of white crappie were significantly different with smaller fish predominating in the diked wetlands. Johnson and others (1997) noted similar findings for white crappies and hypothesized that growth was slower in the diked wetlands. Our data indicate their hypothesis was likely




FIGURE 1. Length frequencies (number per 24-h net day) of white crappies collected from Sandusky Bay and Lake Erie sites. Undiked sites were Muddy Creek Bay and Crane Creek, diked wetlands included OSC (Ottawa Shooting Club), Rusk (Rusk Marshland), and ONWR (Ottawa National Wildlife Refuge). N = total number collected during the 2-4 d sampling period in each wetland. Proportional Stock Density (PSD) is the proportion of fish exceeding 130 mm total length that also exceeded 200 mm total length.

correct. Not only was white crappie age structure different between systems, but growth was significantly less in the diked wetlands. Apparently white crappie populations in diked wetlands can be considered biologically isolated, and we suspect this situation exists for other fish species as well.

Isolation of fish communities in diked wetlands might be expected given that earthen dikes and water control devices create considerable barriers to fish movement into and out of these wetlands. However, the need to remove water in spring and add it in fall may result in a water exchange of up to 75% each year and could provide ample opportunity for fish exchange between diked and undiked areas. We have sampled (unpublished data) the culvert pipe connecting the diked wetland investigated by Johnson and others (1997) to adjacent areas and found white crappies moved both directions during the spring. However, total numbers

leaving were small compared to the white crappie population within the diked wetland and was not extensive enough to prevent Johnson and others (1997) from finding significant differences between the diked wetland and Muddy Creek Bay populations.

Absence of extensive fish exchange between diked wetlands and adjacent undiked areas indicates physical isolation, but we also suspect that differences in ecological conditions between the areas may cause the growth differences we observed. We have unpublished data over several years that suggests larval fish densities in Johnson and others' (1997) study wetland were several orders of magnitude less than in Muddy Creek Bay. White crappie are piscivorous predators when prey fish are available and have been shown to rapidly increase body condition and growth in response to abundant prey (Gabelhouse 1991). This likely accounts for fast growth in undiked areas. Additionally, fish production




FIGURE 2. Age distributions of white crappies collected from Sandusky Bay and Lake Erie sites. Undiked sites were Muddy Creek Bay and Crane Creek, diked wetlands included OSC (Ottawa Shooting Club), Rusk (Rusk Marshland), and ONWR (Ottawa National Wildlife Refuge). N = number of fish that were aged and does not necessarily equal total number collected.




FIGURE 3. Mean backcalculated length-at-age (all year classes combined) of white crappies collected from Sandusky Bay and Lake Erie sites. Undiked sites were Muddy Creek Bay and Crane Creek, diked wetlands included OSC (Ottawa Shooting Club), Rusk (Rusk Marshland), and ONWR (Ottawa National Wildlife Refuge).

may well be limited in diked wetlands by reduced nutrient inputs (Sager and others 1985, Robb and Mitsch 1992), low oxygen and increased water temperatures during summer drawdown (Navarro and Johnson 1992), and the extensive vegetated areas within tying up nutrients. It is our belief that ecological conditions are very different in diked wetlands as compared to adjacent waters and may very well create biologically different fish populations and communities.

White crappie populations in the 3 diked wetlands sampled exhibited considerable differences in the parameters we examined. Rusk and ONWR had white crappie populations that grew significantly slower than the population in OSC. The two former wetlands differed from the latter in two respects; they both 1) contained populations of largemouth bass (*Micropterus salmoides*) and northern pike (*Esox lucius*) whereas the Boardman Unit did not, and 2) undergo less spring drawdown and thus, have higher summer water levels. Adult white crappie must compete for limited prey fish with these two predator species which may limit white crappie growth in Rusk and ONWR. Additionally, carp (*Cyprinus carpio*) densities in these two wetlands were low (0.2 and 0.5 fish per net day during this study), creating clear water and considerable submerged vegetation. White crappie are considered to be a pelagic species and prefers less vegetated areas (Trautman 1981). Therefore, white crappie probably cannot be expected to grow well in vegetated systems.

An often debated subject in Ohio's Lake Erie region relates to management of fish communities in diked wetlands. There are proponents of opening the dikes to enhance fish exchange and allow fish access to possible spawning areas. We presently do not advocate this strategy. We are concerned that "breached" dikes may substantially reduce the present vegetated habitats in diked wetlands crucial to the management of wildlife species, some of which are threatened or endangered. As noted by Johnson and others (1997), research is needed to assess impacts on flora and fauna by allowing continual water exchange. We encourage efforts on improving fish communities in the diked wetlands, focusing on activities such as stocking predators to control carp populations and providing deep water areas for fish communities to use during periods of temperature extremes and low oxygen in shallow areas.

ACKNOWLEDGMENTS. Salaries and support were provided by State and Federal Funds appropriated to the Ohio Agricultural Research and Development Center, The Ohio State University. Manuscript Number 86-97. We thank the Ottawa Shooting Club, the Ottawa National Wildlife Refuge, and Mr. Jerry Rusk for graciously allowing us to sample their wetlands.

## LITERATURE CITED

- Anderson RO, Neumann RM. 1996. Length, weight, and associated structural indices. In: BR Murphy and DW Willis (eds.). Fisheries Techniques, 2nd ed. American Fisheries Society, Bethesda, MD. p. 447-82.
- Bookhout TA, Bednarik KE, Kroll RW. 1989. The Great Lakes Marshes. In: LM Smith, RL Pederson, and RM Kaminski (eds.). Habitat Management for Migrating and Wintering Waterfowl in North America. Lubbock: Texas Tech University Press. p. 131-56.
- Boxrucker J. 1986. A comparison of the otolith and scale methods for aging white crappies in Oklahoma. N. Amer. J. Fish. Mgt. 6: 122-5.
- Everhart WH, Youngs WD. 1975. Principles of Fishery Science. Ithaca, NY: Cornell University Press.
- Gabelhouse DW Jr. 1991. Seasonal changes in body condition of white crappies and relations to length and growth in Melvern Reservoir, Kansas. N. Amer. J. Fish. Mgt. 11: 50-6.
- Herdendorf CE. 1987. The ecology of coastal marshes of western Lake Erie: A community profile. U.S. Fish and Wildlife Service, Office of Biological Services, Washington, DC. Biological Report 85(7.9).
- Hollander M, Wolfe DA. 1973. Nonparametric statistical methods. New York: Wiley.

- Johnson DL. 1989. Lake Erie wetlands: Fisheries considerations. *In*: KA Krieger (ed.). Lake Erie Estuarine Systems: Issues, Resources, Status, and Management. Proceedings of the National Oceanic and Atmospheric Administration Estuary-of-the-Month Seminar Series No. 14, Washington, DC. p. 257-74.
- Johnson DL, Lynch WE Jr., Morrison TW. 1997. Fish communities in a diked Lake Erie wetland and an adjacent undiked area. *Wetlands* 14: 43-54.
- Liston CR, Chubb SL. 1985. Relationships of water level fluctuations and fish. *In*: HH Prince and FM D'Itri (eds.). Coastal Wetlands. Chelsea, MI: Lewis. p. 134-6.
- Navarro JE, Johnson DL. 1992. Ecology of stocked northern pike in

- two Lake Erie controlled wetlands. *Wetlands* 12: 171-7.
- Robb DM, Mitsch WJ. 1992. Selected chemical parameters of diked and undiked Lake Erie marshes. *In*: J Kusler and R Smardon (eds.). Wetlands of the Great Lakes. Assn. of Wetland Managers, Berne, NY. p. 130-36.
- Sager PE, Richman S, Harris HJ, Fewless G. 1985. Preliminary observations on the seiche-induced flux of carbon, nitrogen, and phosphorus in a Great Lakes coastal marsh. *In*: HH Prince and FM D'Itri (eds.). Coastal Wetlands. Chelsea, MI: Lewis. p. 60-8.
- Schramm HL Jr., Doerzbacher JF. 1982. Use of otoliths to age black crappie from Florida. *Proc. Ann. Conf. Southeastern Assoc. Fish and Wildlife Agencies* 36: 95-105.

# The Ohio Academy of Science

## The Ohio Academy of Science

1500 W. Third Ave. Ste. 223  
Columbus, OH  
43212-2817  
Phone or FAX  
(614) 488-2228  
Outside of Franklin Co.  
1-800-OHIOSCI  
oas@iwaynet.net  
http://www.ohiosci.org

Mr. David M. Weaner  
Director, Junior Academy  
Council  
Westerville North HS  
950 County Line Rd.  
Westerville, OH 43081  
(614) 895-6060  
dmweaner@iwaynet.net

Mrs. Connie Hubbard  
Acting Asst Director  
Junior Academy Cncl  
Hoover HS  
525 Seventh St. NE  
N. Canton, OH 44720  
(330) 497-5620  
chubbard@raex.com

## State Science Day Ohio Wesleyan University April 18, 1998

Dr. Michael Grote  
State Science Day Director  
Ohio Wesleyan Univ.  
Dept. of Education  
Delaware, OH 43015  
(614) 368-3561  
mngrote@cc.owu.edu

## District Councilpersons

### District 1 Defiance College March 28, 1998

Dr. Karen McCracken  
Defiance College  
Div. of Science and Math  
Defiance, OH 43512  
(419) 783-2494  
kmcrcracken@tdc.edu


Mr. James Short  
Gorham Fayette HS  
P.O. Box 309  
Fayette, OH 43521  
(419) 237-2114  
Fay\_ACA\_JS@  
nwoca.occn.ohio.gov

### District 2 University of Toledo March 28, 1998

Dr. Mark Camp  
University of Toledo  
Dept. of Geology  
Toledo OH 43606-3390  
(419) 530-2398  
mcamp@geology.utoledo.edu

Ms. Mikell Hedley  
Central Catholic HS  
4531 Woodland Lane  
Sylvania OH 43560  
(419) 885-3031  
mhedley@uof02.utoledo.edu

## 1998 District Science Days



### District 3 OSU Marion March 21, 1998

Dr. Brian W. McEnnis  
OSU-Marion  
1465 Mt. Vernon Ave.  
Marion OH 43302-5695  
(614) 389-6786  
mcennis.1@osu.edu

Ms. Carol White  
Baker MS  
400 Pennsylvania Ave.  
Marion, OH 43302  
(614) 389-6362

### District 4 Ashland University March 28, 1998

Dr. Michael Hudson  
Ashland University  
Ashland, OH 44805  
(419) 289-5270  
mhudson@ashland.edu

Mr. Jeff Steele  
Mapleton HS  
1021 Columbus Circle N.  
Ashland, OH 44805  
(419) 945-2188  
MAPL\_STEELE@tccsa.ohio.gov

### District 5 University of Akron March 28, 1998

Mrs. Toni Miller  
Springfield HS  
721 Northwood Dr.  
Uniontown, OH 44685  
(330) 896-2533  
tmiller@scs.summit.k12.oh.us

### District 6 Ohio Northern Univ. March 28, 1998

Prof. Terry Keiser  
Ohio Northern University  
Dept. of Biological Sciences  
Ada, OH 45810-1599  
(419) 772-2327 or 2325  
t-keiser@onu.edu

Mr. Spencer E. Reames  
Benjamin Logan HS  
209 Blair Ave.  
Bellefontaine, OH 43311  
(937) 592-1666  
sreames@educator.mci.net

### District 7 Columbus State Comm. College March 21, 1998

Dr. Ludwig Sprandel  
Columbus State Comm. College  
Dept. of Biological Sciences  
550 E. Spring St.  
Columbus, OH 43215-1609  
(614) 227-5362  
lsprandel@cscc.edu

Mr. Larry Hohman  
Willard Grizzell MS  
8705 Manley Rd.  
Dublin OH 43017  
(614) 798-3569  
hohmanlr@iwaynet.net

### District 8 OU-Lancaster March 14, 1998

Dr. Steve Nerney  
Ohio University-Lancaster  
1570 Granville Pike  
Lancaster OH 43130-1097  
(614) 654-6711  
nerney@ohiou.edu

Mr. David P. Thrush  
Fairfield Union HS  
6675 Ireland Rd.  
Lancaster OH 43130  
(614) 536-7306

### District 9 Marietta College March 21, 1998

Prof. David Freeman  
Marietta College  
Dept. of Petroleum Engineering  
Marietta, OH 43750  
(614) 376-4778  
freemamd@mcnet.marietta.edu

Mr. Steve Mencer  
Meadowbrook MS  
58607 Marietta Rd.  
Byesville, OH 43723  
(614) 685-2561

### District 10 Wright State Univ. March 28, 1998

Dr. Lois Cook  
Wright State University  
College of Science & Mathematics  
Dayton, OH 45435  
(937) 775-3539

Ms. Phyllis Frysinger  
Miami View School  
230 Old Clifton Rd.  
S. Charleston, OH 45368  
(937) 462-8364  
pfrysing@efanet.com

### District 11 Miami University March 28, 1998

Ms. p. Helen Stevenson  
Miami University  
Physics Dept.  
Oxford, OH 45056  
(513) 529-1859  
stevensonh@muohio.edu

Ms. Donna Davis  
St. Joseph School  
Second & Hanover Streets  
Hamilton OH 45011  
(513) 863-8758

### District 12 Ohio University March 7, 1998

Dr. James Y. Tong  
Ohio University  
Department of Chemistry  
Athens, OH 45701-2979  
(614) 593-1738  
tong@ouvax.cats.ohiou.edu

Mr. Bill Bahr  
River Valley HS  
206 Greer Rd.  
Bidwell, OH 45614-9998  
(614) 446-2642

### District 13 Malone College March 21, 1998

Dr. Ruth Miles  
Malone College  
515 25th St. NW  
Canton OH 44709  
(330) 471-8178  
RMiles@malone.edu

Mrs. Connie Hubbard  
Hoover HS  
525 Seventh St. NE  
N. Canton, OH 44720  
(330) 497-5620  
chubbard@raex.com

### District 14 Shawnee State Univ. March 21, 1998

Ms. Judy Meeker  
Shawnee State University  
Massey Hall B50  
Portsmouth, OH 45662  
(740) 355-2411  
jmeeker@shawnee.edu

Ms. Sherylene Rice  
Clay HS  
44 Clay High St.  
Portsmouth, OH 45662  
(614) 354-6644  
srice@clay.scoea.oh.gov

### District 15 Youngstown State University March 14, 1998

Dr. Jeffrey Dick  
Youngstown State University  
Department of Geology  
Youngstown, OH 44555-0001  
(330) 742-3612  
jdick@cc.ysu.edu

Ms. Madonna Chism  
Community Affairs Rep.  
East Ohio Gas Company  
1165 Rayen Ave.  
Youngstown OH 44502-1394  
(330) 742-8104

### District 16 Belmont Technical College March 28, 1998

Mr. Allen Green  
Belmont Technical College  
120 Fox-Shannon Pl.  
St. Clairsville, OH 43950  
(614) 695-9500  
AGreen@belmont.tcc.oh.us

Mr. Vince Thornburg  
Shadyside HS  
54050 Fulton Hill Rd.  
Bellair OH 43096  
(614) 676-7513